

# Hurwitz Genus Formula

## Unit 22

Gil Cohen

December 30, 2024

# Overview

- 1 Adeles in extensions
- 2 Differentials in extensions
- 3 The co-trace
- 4 Hurwitz Genus Formula

# Hurwitz Genus Formula


Throughout this unit  $F/L$  is a finite separable extension of  $E/K$ .

## Lemma 1

Let  $L/K$  be a finite separable field extension. Let  $V$  be an  $L$ -vector space (and so  $V$  is also a  $K$ -vector space). Let  $T : V \rightarrow K$  be a  $K$ -linear map.

Then,  $\exists! T' : V \rightarrow L$  that is  $L$ -linear s.t.

$$\text{Tr}_{L/K} \circ T' = T.$$



We omit the proof of this fact (see Dan Haran's lecture notes; Chapter 33).

# Adeles - recall

Recall that an **adele** of  $F/L$  is a function  $\alpha : \mathbb{P}(F/L) \rightarrow F$  that maps  $\mathfrak{P} \rightarrow \alpha_{\mathfrak{P}}$  s.t.  $v_{\mathfrak{P}}(\alpha_{\mathfrak{P}}) \geq 0$  almost always.

The set of adeles of  $F/L$  is denoted by  $\mathbb{A}_{F/L}$  or  $\mathbb{A}_F$ . Recall that  $\mathbb{A}_F$  is an  $F$ -algebra. Multiplying by elements of  $F$  is done via the embedding  $F \hookrightarrow \mathbb{A}_F$  where  $x \mapsto [x]$  in which  $[x]_{\mathfrak{P}} = x$ .

For  $\mathfrak{a} \in \mathcal{D}(F/L)$  we defined

$$\Lambda_F(\mathfrak{a}) = \{\alpha \in \mathbb{A}_F \mid \forall \mathfrak{P} \in \mathbb{P}(F/L) \quad v_{\mathfrak{P}}(\alpha) + v_{\mathfrak{P}}(\mathfrak{a}) \geq 0\}.$$

We sometimes write  $\Lambda(\mathfrak{a})$  for short.

$$\mathbb{A}_F \ni \alpha \quad \alpha_{\mathfrak{p}} \quad \begin{array}{l} v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) \geq 0 \\ \text{almost} \\ \text{always} \end{array}$$

$\mathbb{P} \quad \dots \quad \mathfrak{p} \quad \dots$

# Adeles of extensions

## Definition 2


We extend the above definition to extensions.

$$\mathbb{A}_{F/E} = \{\alpha \in \mathbb{A}_F \mid \mathfrak{P}_1 \cap E = \mathfrak{P}_2 \cap E \implies \alpha_{\mathfrak{P}_1} = \alpha_{\mathfrak{P}_2}\}.$$

Note that  $F \hookrightarrow \mathbb{A}_{F/E} \subseteq \mathbb{A}_F$  and so  $\mathbb{A}_{F/E}$  is an  $F$ -subalgebra of  $\mathbb{A}_F$ .

Moreover, for  $\mathfrak{a} \in \mathcal{D}(F/L)$  we define

$$\Lambda_{F/E}(\mathfrak{a}) = \mathbb{A}_{F/E} \cap \Lambda_F(\mathfrak{a}).$$



## Definition 3

We extend  $\text{Tr}_{F/E} : F \rightarrow E$  to the map

$$\text{Tr}_{F/E} : \mathbb{A}_{F/E} \rightarrow \mathbb{A}_E$$

as follows: For  $\alpha \in \mathbb{A}_{F/E}$  and  $\mathfrak{p} \in \mathbb{P}(E)$ ,

$$(\text{Tr}_{F/E}(\alpha))_{\mathfrak{p}} = \text{Tr}_{F/E}(\alpha_{\mathfrak{P}})$$

where  $\mathfrak{P}$  is some prime divisor lying over  $\mathfrak{p}$ .

We need to prove that indeed

$$\text{Tr}_{F/E}(\alpha) \in \mathbb{A}(E).$$

Namely, we need to show that  $\text{Tr}_{F/E}(\alpha)_{\mathfrak{p}} \geq 0$  almost always.

# Adeles of extensions

We need to show that  $\text{Tr}_{F/E}(\alpha)_p \geq 0$  almost always.

As  $\alpha \in \mathbb{A}_{F/E}$  we have that  $\alpha_{\mathfrak{p}} \in \mathcal{O}_{\mathfrak{p}}$  almost always. Thus, for almost all  $\mathfrak{p} \in \mathbb{P}(E)$ ,

$$\forall \mathfrak{p}/\mathfrak{p} \quad \alpha_{\mathfrak{p}} \in \bigcap_{\mathfrak{p}'/\mathfrak{p}} \mathcal{O}_{\mathfrak{p}'} = \mathcal{O}'_{\mathfrak{p}}.$$

Recall that  $\text{Tr}_{F/E}(\mathcal{O}'_{\mathfrak{p}}) = \mathcal{O}_{\mathfrak{p}}$ , and so for almost all  $\mathfrak{p}$ ,

$$(\text{Tr}_{F/E}(\alpha))_{\mathfrak{p}} = \text{Tr}_{F/E}(\alpha_{\mathfrak{p}}) \in \mathcal{O}_{\mathfrak{p}},$$

thus establishing that  $\text{Tr}_{F/E}(\alpha) \in \mathbb{A}(E)$ .

# Adeles of extensions

We further remark that

$$\mathrm{Tr}_{F/E}([x]) = [\mathrm{Tr}_{F/E}(x)].$$

## Lemma 4

For every  $\mathfrak{a} \in \mathcal{D}(F)$  we have that

$$\mathbb{A}_F = \mathbb{A}_{F/E} + \Lambda_F(\mathfrak{a}).$$

## Proof.

The inclusion  $\mathbb{A}_F \supset \mathbb{A}_{F/E} + \Lambda_F(\mathfrak{a})$  is obvious. For the other inclusion, take  $\alpha \in \mathbb{A}_F$ . We first construct some  $\beta \in \mathbb{A}_{F/E}$  as follows.



# Adeles of extensions

Proof.

Take  $\mathfrak{p} \in \mathbb{P}(E)$ . The set of  $\mathfrak{P}/\mathfrak{p}$  is finite and so by WAT,  $\exists x_{\mathfrak{p}} \in F$  s.t.

$$\forall \mathfrak{P}/\mathfrak{p} \quad v_{\mathfrak{P}}(x_{\mathfrak{p}} - \alpha_{\mathfrak{P}}) \geq -v_{\mathfrak{P}}(\mathfrak{a}).$$

Note that for almost all  $\mathfrak{p} \in \mathbb{P}(E)$  we have that  $\forall \mathfrak{P}/\mathfrak{p} \quad v_{\mathfrak{P}}(\mathfrak{a}) = 0$ .

Moreover, since  $\alpha \in \mathbb{A}_F$ , for almost all  $\mathfrak{P} \in \mathbb{P}(F)$ ,  $v_{\mathfrak{P}}(\alpha_{\mathfrak{P}}) \geq 0$ . Thus, for almost all  $\mathfrak{p}$ ,

$$\begin{aligned} \forall \mathfrak{P}/\mathfrak{p} \quad v_{\mathfrak{P}}(x_{\mathfrak{p}}) &= v_{\mathfrak{P}}(x_{\mathfrak{p}} - \alpha_{\mathfrak{P}} + \alpha_{\mathfrak{P}}) \\ &\geq \min(v_{\mathfrak{P}}(x_{\mathfrak{p}} - \alpha_{\mathfrak{P}}), v_{\mathfrak{P}}(\alpha_{\mathfrak{P}})) \\ &\geq 0. \end{aligned}$$

## Proof.

With this, we define  $\beta : \mathbb{P}(F) \rightarrow F$  by

$$\beta_{\mathfrak{P}} = x_{\mathfrak{p}},$$

where  $\mathfrak{p} \in \mathbb{P}(E)$  is the prime divisor lying under  $\mathfrak{P}$ .

$\beta \in \mathbb{A}_F$  as  $v_{\mathfrak{P}}(\beta_{\mathfrak{P}}) = v_{\mathfrak{P}}(x_{\mathfrak{p}}) \geq 0$  almost always. Moreover,  $\beta \in \mathbb{A}(F/E)$  since we take  $\beta_{\mathfrak{P}} = x_{\mathfrak{p}} = \beta_{\mathfrak{P}'}$  for all places  $\mathfrak{P}, \mathfrak{P}'$  lying over  $\mathfrak{p}$ .

Lastly, note that  $\alpha - \beta \in \Lambda_F(\mathfrak{a})$ . Indeed,  $\forall \mathfrak{P} \in \mathbb{P}(F)$ ,

$$v_{\mathfrak{P}}(\alpha - \beta) = v_{\mathfrak{P}}(\alpha_{\mathfrak{P}} - \beta_{\mathfrak{P}}) = v_{\mathfrak{P}}(\alpha_{\mathfrak{P}} - x_{\mathfrak{p}}) \geq -v_{\mathfrak{P}}(\mathfrak{a}).$$

Thus,

$$\alpha = \beta + (\alpha - \beta) \in \mathbb{A}_{F/E} + \Lambda_F(\mathfrak{a}),$$

concluding the proof. □

# Overview

- 1 Adeles in extensions
- 2 Differentials in extensions
- 3 The co-trace
- 4 Hurwitz Genus Formula

# Differential - recall

Recall that a **differential** of  $F/L$  is an  $L$ -linear map  $\omega : \mathbb{A}_F \rightarrow L$  that is nullified on a subspace of the form  $\Lambda(\mathfrak{a}) + F$  for some divisor  $\mathfrak{a}$ .

$$\omega : \begin{array}{c} \mathbb{A}_{F/L} \\ \text{---} \\ \alpha : \mathbb{P}_{F/L} \rightarrow F \end{array} \longrightarrow L$$

For a differential  $\omega \neq 0$  we defined the canonical divisor

$$(\omega) = \max \{ \mathfrak{a} \in \mathcal{D}(F) : \omega|_{\Lambda(\mathfrak{a})+F} = 0 \}.$$

In particular,  $\omega|_{\Lambda((\omega))} = 0$ .

# Differentials in extensions

## Lemma 5

Let  $\omega : \mathbb{A}_E \rightarrow K$  be a differential of  $E/K$ . We define a map

$$\omega_1 : \mathbb{A}_{F/E} \rightarrow K$$

by  $\omega_1 = \omega \circ \text{Tr}_{F/E}$ . Then,

- 1  $\omega_1$  is  $K$ -linear; and
- 2  $\omega_1$  is nullified on  $\Lambda_{F/E}(\mathfrak{a}) + F$ , where

$$\mathfrak{a} = \text{Con}_{F/E}(\omega) + \text{Diff}(F/E).$$

A commutative diagram with  $\mathbb{A}_{F/E}$  at the top left,  $K$  at the top right, and  $\mathbb{A}_E$  at the bottom left. A horizontal arrow labeled  $\omega_1$  points from  $\mathbb{A}_{F/E}$  to  $K$ . A vertical arrow labeled  $\text{Tr}_{F/E}$  points from  $\mathbb{A}_{F/E}$  down to  $\mathbb{A}_E$ . A diagonal arrow labeled  $\omega$  points from  $\mathbb{A}_E$  up to  $K$ .

# Differentials in extensions

## Proof.

The first item follows since both  $\text{Tr}_{F/E}$  and  $\omega$  are  $K$ -linear maps.

For the second item, first note that  $\omega_1|_F = 0$ . Indeed,  $\text{Tr}_{F/E}(F) = E$ , and  $\omega|_E = 0$ .

We turn to prove that  $(\omega_1)|_{\Lambda_{F/E}(\mathfrak{a})} = 0$ .

Take  $\alpha \in \Lambda_{F/E}(\mathfrak{a})$ . We need to show that  $\omega_1(\alpha) = \omega(\text{Tr}_{F/E}(\alpha)) = 0$ . To this end we show that

$$\text{Tr}_{F/E}(\alpha) \in \Lambda_E((\omega)).$$

Equivalently,

$$\forall \mathfrak{p} \in \mathbb{P}(E) \quad v_{\mathfrak{p}}(\text{Tr}_{F/E}(\alpha)) + v_{\mathfrak{p}}((\omega)) \geq 0.$$

Thus, we need to show that for all  $\mathfrak{p}$  and  $\mathfrak{A}/\mathfrak{p}$ ,

$$v_{\mathfrak{p}}(\text{Tr}_{F/E}(\alpha_{\mathfrak{A}})) + v_{\mathfrak{p}}((\omega)) \geq 0.$$

# Differentials in extensions

Proof.

We want to show that

$$\forall \mathfrak{p}, \mathfrak{P}/\mathfrak{p} \quad v_{\mathfrak{p}}(\mathrm{Tr}_{F/E}(\alpha_{\mathfrak{P}})) + v_{\mathfrak{p}}((\omega)) \geq 0.$$

Fix  $\mathfrak{p}$  and let  $x \in E$  be s.t.  $v_{\mathfrak{p}}(x) = v_{\mathfrak{p}}((\omega))$ . Then, for all  $\mathfrak{P}/\mathfrak{p}$ ,

$$\begin{aligned} v_{\mathfrak{P}}(x\alpha_{\mathfrak{P}}) &= v_{\mathfrak{P}}(x) + v_{\mathfrak{P}}(\alpha_{\mathfrak{P}}) \\ &= e(\mathfrak{P}/\mathfrak{p})v_{\mathfrak{p}}(x) + v_{\mathfrak{P}}(\alpha_{\mathfrak{P}}) \\ &= e(\mathfrak{P}/\mathfrak{p})v_{\mathfrak{p}}((\omega)) + v_{\mathfrak{P}}(\alpha_{\mathfrak{P}}) \\ &\geq e(\mathfrak{P}/\mathfrak{p})v_{\mathfrak{p}}((\omega)) - v_{\mathfrak{P}}(\mathfrak{a}) \\ &= v_{\mathfrak{P}}(\mathrm{Con}_{F/E}((\omega)) - \mathfrak{a}) \\ &= v_{\mathfrak{P}}(-\mathrm{Diff}(F/E)) \\ &= -d(\mathfrak{P}/\mathfrak{p}). \end{aligned}$$

Thus,  $x\alpha_{\mathfrak{P}} \in \mathcal{C}_{\mathfrak{p}}$ .

Proof.

Fix  $\mathfrak{p}$  and let  $x \in E$  be s.t.  $v_{\mathfrak{p}}(x) = v_{\mathfrak{p}}((\omega))$ . Then,  $x\alpha_{\mathfrak{p}} \in \mathcal{C}_{\mathfrak{p}}$ . Thus,

$$v_{\mathfrak{p}}(\mathrm{Tr}_{F/E}(x\alpha_{\mathfrak{p}})) \geq 0.$$

Since  $\mathrm{Tr}_{F/E}$  is  $E$ -linear, we get that

$$\begin{aligned} v_{\mathfrak{p}}(\mathrm{Tr}_{F/E}(x\alpha_{\mathfrak{p}})) &= v_{\mathfrak{p}}(x\mathrm{Tr}_{F/E}(\alpha_{\mathfrak{p}})) \\ &= v_{\mathfrak{p}}(x) + v_{\mathfrak{p}}(\mathrm{Tr}_{F/E}(\alpha_{\mathfrak{p}})) \\ &= v_{\mathfrak{p}}((\omega)) + v_{\mathfrak{p}}(\mathrm{Tr}_{F/E}(\alpha_{\mathfrak{p}})). \end{aligned}$$

Thus,

$$v_{\mathfrak{p}}((\omega)) + v_{\mathfrak{p}}(\mathrm{Tr}_{F/E}(\alpha_{\mathfrak{p}})) \geq 0$$

which, recall, concludes the proof. □



# Differentials in extensions

Let  $\omega : \mathbb{A}_E \rightarrow K$  be a differential of  $E/K$ . Recall that we have defined the map

$$\omega_1 : \mathbb{A}_{F/E} \rightarrow K$$

by  $\omega_1 = \omega \circ \text{Tr}_{F/E}$ . We further denoted

$$\mathfrak{a} = \text{Con}_{F/E}(\omega) + \text{Diff}(F/E).$$

and proved that  $\omega_1$  is nullified on  $F + \Lambda_{F/E}(\mathfrak{a})$ .

## Lemma 6

$\forall \mathfrak{a}' \in \mathcal{D}(F)$

$$\mathfrak{a}' \not\subseteq \mathfrak{a} \implies (\omega_1)|_{\Lambda_{F/E}(\mathfrak{a}')} \neq 0.$$

# Differentials in extensions

Proof. (Proof of Lemma 6)

We wish to prove that

$$\forall \alpha' \in \mathcal{D}(F) \text{ s.t. } \alpha' \not\leq \alpha \quad \exists \beta \in \Lambda_{F/E}(\alpha') \text{ s.t. } \omega_1(\beta) \neq 0.$$

Fix  $\alpha' \not\leq \alpha$  and let  $\mathfrak{P}' \in \mathbb{P}(F)$  s.t.

$$v_{\mathfrak{P}'}(\alpha') > v_{\mathfrak{P}'}(\alpha) = v_{\mathfrak{P}'}(\text{Con}_{F/E}(\omega)) + d(\mathfrak{P}'/\mathfrak{p}),$$

where  $\mathfrak{p}$  is the prime divisor lying under  $\mathfrak{P}'$ . That is,

$$v_{\mathfrak{P}'}(\text{Con}_{F/E}(\omega) - \alpha') < -d(\mathfrak{P}'/\mathfrak{p}).$$

Define

$$J = \{z \in F \mid \forall \mathfrak{P}/\mathfrak{p} \quad v_{\mathfrak{P}}(z) \geq v_{\mathfrak{P}}(\text{Con}_{F/E}(\omega) - \alpha')\}.$$

$J$  is closed under addition and under multiplication by  $\mathcal{O}'_{\mathfrak{p}}$  and so  $J$  is an  $\mathcal{O}'_{\mathfrak{p}}$ -module. Furthermore,  $\text{Tr}_{F/E}(J)$  is an  $\mathcal{O}_{\mathfrak{p}}$ -module.

Proof.

$$J = \{z \in F \mid \forall \mathfrak{P}/\mathfrak{p} \ v_{\mathfrak{P}}(z) \geq v_{\mathfrak{P}}(\text{Con}_{F/E}(\omega) - \mathfrak{a}')\}.$$

By WAT  $\exists z' \in F$  s.t.

$$\forall \mathfrak{P}/\mathfrak{p} \ v_{\mathfrak{P}}(z') = v_{\mathfrak{P}}(\text{Con}_{F/E}(\omega) - \mathfrak{a}').$$

In particular,  $z' \in J$  and

$$v_{\mathfrak{P}'}(z') < -d(\mathfrak{P}'/\mathfrak{p}),$$

and so  $z' \notin C_{\mathfrak{p}}$ . Thus,  $\exists v \in \mathcal{O}'_{\mathfrak{p}}$  s.t.

$$\text{Tr}_{F/E}(vz') \notin \mathcal{O}_{\mathfrak{p}}.$$

As  $J$  is an  $\mathcal{O}'_{\mathfrak{p}}$ -module,  $vz' \in J$  and so  $\text{Tr}_{F/E}(J) \not\subseteq \mathcal{O}_{\mathfrak{p}}$ .

# Differentials in extensions

Proof.

$$J = \{z \in F \mid \forall \mathfrak{P}/\mathfrak{p} \ v_{\mathfrak{P}}(z) \geq v_{\mathfrak{P}}(\text{Con}_{F/E}(\omega) - \mathfrak{a}')\}.$$

Let  $t \in E$  be with  $v_{\mathfrak{p}}(t) = 1$ . Thus, for a sufficiently large  $r$ ,

$$t^r J \subseteq \bigcap_{\mathfrak{P}/\mathfrak{p}} \mathcal{O}_{\mathfrak{P}} = \mathcal{O}'_{\mathfrak{p}}.$$

Hence,

$$t^r \text{Tr}_{F/E}(J) = \text{Tr}_{F/E}(t^r J) \subseteq \mathcal{O}_{\mathfrak{p}} \implies v_{\mathfrak{p}}(\text{Tr}_{F/E}(J)) \geq -r.$$

In this case, we proved in a previous unit that

$$\text{Tr}_{F/E}(J) = t^m \mathcal{O}_{\mathfrak{p}}$$

for some  $m \in \mathbb{Z}$ . In our case  $m \leq -1$  as otherwise  $\text{Tr}_{F/E}(J) \subseteq \mathcal{O}_{\mathfrak{p}}$ .

# Differentials in extensions

Proof.

Recall that  $(\omega)$  is the largest divisor in  $\mathcal{D}(E)$  on which  $\omega$  vanishes. Thus,  $\omega$  does not vanish on  $\Lambda_E((\omega) + \mathfrak{p})$ . Namely,

$$\exists \alpha \in \Lambda_E((\omega) + \mathfrak{p}) \quad \text{s.t.} \quad \omega(\alpha) \neq 0.$$

Note that  $\alpha \notin \Lambda_E((\omega))$ .

Since for all other prime divisors  $\mathfrak{q} \neq \mathfrak{p}$  we have

$$v_{\mathfrak{q}}((\omega)) = v_{\mathfrak{q}}((\omega) + \mathfrak{p})$$

we conclude that

$$\begin{aligned} v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) &\geq -v_{\mathfrak{p}}((\omega) + \mathfrak{p}) = -v_{\mathfrak{p}}((\omega)) - 1, \\ v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) &\not\geq -v_{\mathfrak{p}}((\omega)), \end{aligned}$$

and so

$$v_{\mathfrak{p}}(\alpha_{\mathfrak{p}}) = -v_{\mathfrak{p}}((\omega)) - 1.$$

# Differentials in extensions

Proof.

Define  $\gamma, \gamma' \in \mathbb{A}_E$  as follows

$$\gamma_{\mathfrak{q}} = \begin{cases} \alpha_{\mathfrak{p}}, & \mathfrak{q} = \mathfrak{p} \\ 0, & \mathfrak{q} \neq \mathfrak{p}. \end{cases} \quad \gamma'_{\mathfrak{q}} = \begin{cases} 0, & \mathfrak{q} = \mathfrak{p} \\ \alpha_{\mathfrak{q}}, & \mathfrak{q} \neq \mathfrak{p}. \end{cases}$$

Note that

- 1  $\gamma, \gamma'$  are adeles;
- 2  $\gamma + \gamma' = \alpha$ ;
- 3  $\gamma' \in \Lambda_E((\omega))$ ; and so  $\omega(\gamma') = 0$ ;
- 4  $\omega(\gamma) = \omega(\alpha) - \omega(\gamma') = \omega(\alpha) \neq 0$ .

Write  $x = \gamma_{\mathfrak{p}} = \alpha_{\mathfrak{p}}$ . Take  $y \in E$  s.t.  $v_{\mathfrak{p}}(y) = v_{\mathfrak{p}}((\omega))$ . Then,

$$v_{\mathfrak{p}}(xy) = v_{\mathfrak{p}}(x) + v_{\mathfrak{p}}(y) = (-v_{\mathfrak{p}}((\omega)) - 1) + v_{\mathfrak{p}}((\omega)) = -1 \geq m.$$

Hence,  $xy \in t^m \mathcal{O}_{\mathfrak{p}}$ .

# Differentials in extensions

Proof.

Recall that  $\text{Tr}_{F/E}(J) = t^m \mathcal{O}_{\mathfrak{p}}$  and  $xy \in t^m \mathcal{O}_{\mathfrak{p}}$ , and so

$$\exists z \in J \text{ s.t. } \text{Tr}_{F/E}(z) = xy.$$

Define an adèle  $\beta \in \mathbb{A}_{F/E}$  by

$$\beta_{\mathfrak{p}} = \begin{cases} zy^{-1}, & \mathfrak{p}/\mathfrak{p}; \\ 0, & \text{otherwise.} \end{cases}$$

As  $z \in J$  we have that

$$\forall \mathfrak{p}/\mathfrak{p} \quad v_{\mathfrak{p}}(z) \geq v_{\mathfrak{p}}(\text{Con}_{F/E}(\omega) - \mathfrak{a}').$$

Thus, using that  $v_{\mathfrak{p}}(y) = v_{\mathfrak{p}}((\omega))$

$$\begin{aligned} v_{\mathfrak{p}}(\beta) &= v_{\mathfrak{p}}(z) - v_{\mathfrak{p}}(y) \\ &\geq v_{\mathfrak{p}}(\text{Con}_{F/E}(\omega) - \mathfrak{a}') - v_{\mathfrak{p}}(\text{Con}_{F/E}((\omega))) = -v_{\mathfrak{p}}(\mathfrak{a}'). \end{aligned}$$

Proof.

For  $\mathfrak{P}$  not over  $\mathfrak{p}$ ,

$$v_{\mathfrak{P}}(\beta) = v_{\mathfrak{P}}(0) = \infty > -v_{\mathfrak{P}}(\mathfrak{a}'),$$

and so  $\beta \in \Lambda_{F/E}(\mathfrak{a}')$ . Next, we show that  $\mathrm{Tr}_{F/E}(\beta) = \gamma$ . Indeed,

$$\mathrm{Tr}_{F/E}(\beta)_{\mathfrak{p}} = \mathrm{Tr}_{F/E}(zy^{-1}) = y^{-1}\mathrm{Tr}_{F/E}(z) = y^{-1}yx = \gamma_{\mathfrak{p}}.$$

For  $\mathfrak{q} \neq \mathfrak{p}$ ,

$$\mathrm{Tr}_{F/E}(\beta)_{\mathfrak{q}} = \mathrm{Tr}_{F/E}(0) = 0 = \gamma_{\mathfrak{q}}.$$

Thus,

$$\omega_1(\beta) = \omega(\mathrm{Tr}_{F/E}(\beta)) = \omega(\gamma) \neq 0.$$





# Differentials in extensions


## Theorem 7

For every differential  $\omega$  of  $E/K \exists!$  differential  $\omega'$  of  $F/L$  s.t.

$$\forall \beta \in \mathbb{A}_{F/E} \quad \text{Tr}_{L/K}(\omega'(\beta)) = \omega(\text{Tr}_{F/E}(\beta)).$$

Furthermore, if  $\omega \neq 0$  then  $\omega' \neq 0$  and

$$(\omega') = \text{Con}_{F/E}(\omega) + \text{Diff}(F/E).$$



# Differentials in extensions


Proof.

Set

$$\mathfrak{a} = \text{Con}_{F/E}(\omega) + \text{Diff}(F/E).$$

Define  $\omega_1 : \mathbb{A}_{F/E} \rightarrow K$  by

$$\omega_1 = \omega \circ \text{Tr}_{F/E}.$$



# Differentials in extensions


Proof.

Recall Lemma 4 which stated that

$$\forall b \in \mathcal{D}(F/L) \quad \mathbb{A}_F = \mathbb{A}_{F/E} + \Lambda_F(b).$$

Using this we will extend  $\omega_1$  to  $\omega_2 : \mathbb{A}_F \rightarrow K$  as follows: Every element of  $\mathbb{A}_F$  can be written as  $\beta + \gamma$  where  $\beta \in \mathbb{A}_{F/E}$  and  $\gamma \in \Lambda_F(\alpha)$ . We define

$$\omega_2(\beta + \gamma) = \omega_1(\beta).$$



# Differential

Proof.


Every element of  $\mathbb{A}_F$  can be written as  $\beta + \gamma$  where  $\beta \in \mathbb{A}_{F/E}$  and  $\gamma \in \Lambda_F(\mathfrak{a})$ . We define

$$\omega_2(\beta + \gamma) = \omega_1(\beta).$$

Note that taking  $\gamma = 0 \in \Lambda_F(\mathfrak{a})$  we get

$$\omega_2(\beta) = \omega_2(\beta + 0) = \omega_1(\beta),$$

and so  $\omega_2$  does indeed extend  $\omega_1$ .



Proof.

We turn to show that  $\omega_2$  is well defined.

If  $\beta_1 + \gamma_1 = \beta_2 + \gamma_2$  then

$$\beta_1 - \beta_2 = \gamma_2 - \gamma_1 \in \mathbb{A}_{F/E} \cap \Lambda_F(\mathfrak{a}) = \Lambda_{F/E}(\mathfrak{a}).$$

By Lemma 5,  $\omega_1$  is nullified on  $\Lambda_{F/E}(\mathfrak{a}) + F$  and so

$$\omega_1(\beta_1) - \omega_1(\beta_2) = \omega_1(\beta_1 - \beta_2) = 0.$$

Therefore,

$$\omega_2(\beta_1 + \gamma_1) = \omega_1(\beta_1) = \omega_1(\beta_2) = \omega_2(\beta_2 + \gamma_2).$$

Hence,  $\omega_2$  is well-defined.

# Differentials in extensions


Proof.

Since  $\omega_1$  is  $K$ -linear so is  $\omega_2$ . Lemma 1 then implies that

$$\exists! \omega' : \mathbb{A}_F \rightarrow L \quad \text{s.t.} \quad \text{Tr}_{L/K} \circ \omega' = \omega_2.$$

We want to show that

$$\forall \beta \in \mathbb{A}_{F/E} \quad \text{Tr}_{L/K}(\omega'(\beta)) = \omega(\text{Tr}_{F/E}(\beta)).$$



# Differentials in extensions


Proof.

We want to show that

$$\forall \beta \in \mathbb{A}_{F/E} \quad \text{Tr}_{L/K}(\omega'(\beta)) = \omega(\text{Tr}_{F/E}(\beta)).$$

For every  $\beta \in \mathbb{A}_{F/E}$  we have

$$\text{Tr}_{L/K}(\omega'(\beta)) = \omega_2(\beta) = \omega_1(\beta) = \omega(\text{Tr}_{F/E}(\beta)).$$



# Differentials in extensions

Proof.


We turn to prove that  $\omega'$  is a differential. To this end, we will show that  $\omega'$  vanishes on  $\Lambda_F(\mathfrak{a}) + F$ .

Otherwise, since  $\omega' : \mathbb{A}_F \rightarrow L$  is  $L$ -linear we will have that

$$\omega'(\Lambda_F(\mathfrak{a}) + F) = L.$$

As  $\text{Tr}_{L/K}$  is onto  $K$ , we have that

$$\text{Tr}_{L/K}(\omega'(\Lambda_F(\mathfrak{a}) + F)) = K \implies \omega_2(\Lambda_F(\mathfrak{a}) + F) = K.$$





# Differentials in extensions

Proof.

$$\omega_2(\Lambda_F(\mathfrak{a}) + F) = K. \quad (1)$$

Recall that every element of  $\mathbb{A}_F$  can be written as  $\beta + \gamma$  where  $\beta \in \mathbb{A}_{F/E}$  and  $\gamma \in \Lambda_F(\mathfrak{a})$ , and that we defined

$$\omega_2(\beta + \gamma) = \omega_1(\beta).$$

Thus,

$$\omega_2(\Lambda_F(\mathfrak{a})) = \omega_1(0) = 0. \quad (2)$$

Further, by Lemma 5,  $\omega_1(F) = 0$ . Since  $F \hookrightarrow \mathbb{A}_{F/E}$  and  $\omega_2$  extend  $\omega_1$  on  $\mathbb{A}_{F/E}$  we have that

$$\omega_2(F) = 0. \quad (3)$$

Equations (2),(3) imply

$$\omega_2(\Lambda_F(\mathfrak{a}) + F) = 0,$$

in contradiction to Equation (1).



# Differentials in extensions


Proof.

We turn to establish uniqueness. Take a differential  $\omega'' : \mathbb{A}_F \rightarrow L$  s.t.

$$\forall \beta \in \mathbb{A}_{F/E} \quad \text{Tr}_{L/K}(\omega''(\beta)) = \text{Tr}_{L/K}(\omega'(\beta)) = \omega(\text{Tr}_{F/E}(\beta)).$$

Then,  $\eta = \omega'' - \omega'$  is a differential of  $F/L$  and, in particular is  $L$ -linear, so

$$\forall \beta \in \mathbb{A}_{F/E} \quad \text{Tr}_{L/K}(\eta(\beta)) = \text{Tr}_{L/K}(\omega''(\beta)) - \text{Tr}_{L/K}(\omega'(\beta)) = 0.$$



# Differentials in extensions

Proof.


$$\mathrm{Tr}_{L/K}(\eta(\mathbb{A}_{F/E})) = 0.$$

As  $\mathrm{Tr}_{L/K}$  is onto, we have that

$$\eta(\mathbb{A}_{F/E}) \not\subseteq L.$$

By the L-linearity of  $\eta$ , we get that  $\eta(\mathbb{A}_{F/E}) = 0$ .

Since  $\eta$  is a differential it also vanishes on some  $\Lambda_F(\mathfrak{b})$  for some divisor  $\mathfrak{b}$  and so, by Lemma 4,  $\eta$  vanishes on  $\mathbb{A}_F$ , namely,  $\omega' = \omega''$ .



# Differentials in extensions

Proof.


To conclude the proof, we show that

$$(\omega') = \mathfrak{a} = \text{Con}_{F/E}(\omega) + \text{Diff}(F/E).$$

We already proved that  $\omega'$  vanishes on  $\mathfrak{a}$ , and so we need to prove that  $\mathfrak{a}$  is the largest such divisor.

To this end, take  $\mathfrak{a}' \in \mathcal{D}(F)$  s.t.  $\mathfrak{a}' \not\subseteq \mathfrak{a}$ . We will show that

$$\exists \beta \in \Lambda_F(\mathfrak{a}') \quad \text{s.t.} \quad \omega'(\beta) \neq 0.$$



# Differentials in extensions

Proof.

We wish to show that


$$\mathfrak{a}' \not\subseteq \mathfrak{a} \implies \exists \beta \in \Lambda_F(\mathfrak{a}') \text{ s.t. } \omega'(\beta) \neq 0.$$

By Lemma 6,

$$\exists \beta \in \Lambda_{F/E}(\mathfrak{a}') \subseteq \Lambda_F(\mathfrak{a}') \text{ s.t. } \omega_1(\beta) \neq 0.$$

However,  $\beta \in \Lambda_{F/E}(\mathfrak{a}')$  and so  $\omega_2(\beta) = \omega_1(\beta) \neq 0$ .

As  $\omega_2(\beta) = \text{Tr}_{L/K}(\omega'(\beta))$  we conclude that  $\omega'(\beta) \neq 0$ .



# Overview

- 1 Adeles in extensions
- 2 Differentials in extensions
- 3 The co-trace
- 4 Hurwitz Genus Formula

# The co-trace

## Definition 8


The map

$$\begin{aligned} \text{cotr}_{F/E} : \Omega_{E/K} &\rightarrow \Omega_{F/L} \\ \omega &\mapsto \omega' \end{aligned}$$

that is defined implicitly by the property

$$\text{Tr}_{L/K} \circ \text{cotr}_{F/E}(\omega) = \omega \circ \text{Tr}_{F/E}$$

on  $\mathbb{A}_{F/E}$  is called the **co-trace**.



## Claim 9

Let  $\omega_1, \omega_2 \in \Omega_{E/K}$ . Then,

$$\text{cotr}_{F/E}(\omega_1 + \omega_2) = \text{cotr}_{F/E}(\omega_1) + \text{cotr}_{F/E}(\omega_2).$$

## Proof.

We have that

$$\text{Tr}_{L/K} \circ \text{cotr}_{F/E}(\omega_1) = \omega_1 \circ \text{Tr}_{F/E},$$

$$\text{Tr}_{L/K} \circ \text{cotr}_{F/E}(\omega_2) = \omega_2 \circ \text{Tr}_{F/E}.$$

Thus,

$$\begin{aligned} \text{Tr}_{L/K} \circ (\text{cotr}_{F/E}(\omega_1) + \text{cotr}_{F/E}(\omega_2)) &= \\ \text{Tr}_{L/K} \circ \text{cotr}_{F/E}(\omega_1) + \text{Tr}_{L/K} \circ \text{cotr}_{F/E}(\omega_2) &= \\ \omega_1 \circ \text{Tr}_{F/E} + \omega_2 \circ \text{Tr}_{F/E} &= (\omega_1 + \omega_2) \circ \text{Tr}_{F/E}, \end{aligned}$$

and the proof follows by the (implicit) definition of  $\text{cotr}_{F/E}(\omega_1 + \omega_2)$ .



# The co-trace

Recall that for  $\omega \in \Omega_{E/K}$  and  $x \in E$ , we defined  $x\omega \in \Omega_{E/K}$  by

$$\forall \alpha \in \mathbb{A}_E \quad (x\omega)(\alpha) = \omega(x\alpha).$$

## Claim 10

Let  $\omega \in \Omega_{E/K}$ , and  $x \in E$ . Then,

$$\text{cotr}_{F/E}(x\omega) = x \cdot \text{cotr}_{F/E}(\omega).$$

## Proof.

Let

$$\begin{aligned} \varphi_x : \mathbb{A}_{F/E} &\rightarrow \mathbb{A}_{F/E} \\ \alpha &\mapsto x\alpha. \end{aligned}$$

Recall that by the implicit definition of  $\text{cotr}_{F/E}$  we have that on  $\mathbb{A}_{F/E}$ ,

$$\text{Tr}_{L/K} \circ \text{cotr}_{F/E}(\omega) = \omega \circ \text{Tr}_{F/E}.$$

Proof.

Thus,

$$\mathrm{Tr}_{L/K} \circ \mathrm{cotr}_{F/E}(\omega) \circ \varphi_x = \omega \circ \mathrm{Tr}_{F/E} \circ \varphi_x.$$

Now, for every  $\alpha \in \mathbb{A}_{F/E}$ ,

$$(\mathrm{Tr}_{F/E} \circ \varphi_x)(\alpha) = \mathrm{Tr}_{F/E}(x\alpha) = x\mathrm{Tr}_{F/E}(\alpha) = (\varphi_x \circ \mathrm{Tr}_{F/E})(\alpha).$$

Therefore, on  $\mathbb{A}_{F/E}$ ,

$$\mathrm{Tr}_{F/E} \circ \varphi_x = \varphi_x \circ \mathrm{Tr}_{F/E}.$$

Thus, on  $\mathbb{A}_{F/E}$ ,

$$\mathrm{Tr}_{L/K} \circ \mathrm{cotr}_{F/E}(\omega) \circ \varphi_x = \omega \circ \varphi_x \circ \mathrm{Tr}_{F/E}.$$

Proof.

$$\mathrm{Tr}_{L/K} \circ \mathrm{cotr}_{F/E}(\omega) \circ \varphi_x = \omega \circ \varphi_x \circ \mathrm{Tr}_{F/E}.$$

But

$$x\omega = \omega \circ \varphi_x,$$

$$x \cdot \mathrm{cotr}_{F/E}(\omega) = \mathrm{cotr}_{F/E}(\omega) \circ \varphi_x,$$

and so on  $\mathbb{A}_{F/E}$ ,

$$\mathrm{Tr}_{L/K} \circ (x \cdot \mathrm{cotr}_{F/E}(\omega)) = (x\omega) \circ \mathrm{Tr}_{F/E}.$$

The proof follows by the (implicit) definition of  $\mathrm{cotr}_{F/E}$ . □

## Claim 11

Let  $F/E$  and  $F'/F$  be finite separable extensions of function fields. Then,

$$\text{cotr}_{F'/E} = \text{cotr}_{F'/F} \circ \text{cotr}_{F/E}.$$

As with all tower type statement, we omit the proof.

# Overview

- 1 Adeles in extensions
- 2 Differentials in extensions
- 3 The co-trace
- 4 Hurwitz Genus Formula**

## Theorem 12

Let  $F/L$  be a finite separable extension of  $E/K$ . Let  $g_E, g_F$  be the corresponding genera. Then,

$$2g_F - 2 = \frac{[F : E]}{[L : K]} \cdot (2g_E - 2) + \deg \text{Diff}(F/E).$$

# Hurwitz Genus Formula

$$2g_F - 2 = \frac{[F : E]}{[L : K]} \cdot (2g_E - 2) + \deg \text{Diff}(F/E).$$

Proof.

Take  $0 \neq \omega \in \Omega_{E/K}$ . By Theorem 7,

$$(\text{cotr}_{F/E}(\omega)) = \text{Con}_{F/E}((\omega)) + \text{Diff}(F/E). \quad (4)$$

As  $(\omega)$ ,  $(\text{cotr}_{F/E}(\omega))$  are canonical divisors of  $E/K$  and  $F/L$ , respectively, Riemann-Roch theorem implies that

$$\deg_E((\omega)) = 2g_E - 2 \quad \deg_F((\text{cotr}_{F/E}(\omega))) = 2g_F - 2.$$

The proof then follows by taking  $\deg_F$  on Equation (4) and using that

$$\deg_F(\text{Con}_{F/E}((\omega))) = \frac{[F : E]}{[L : K]} \cdot \deg_E((\omega)).$$

